

# Serotonin rabbit pAb – IS1007

Ref: IS1007-sp

The anti-Serotonin antibody IS1007 is a rabbit polyclonal antibody with high affinity and specificity. Using with the [STAINperfect immunostaining kit A](#), our antibody labeled directly 5-HT in whole mounts, cell culture and tissue sections.

<b>Clonality</b>	Polyclonal antibody
<b>Host</b>	Rabbit
<b>Reactivity</b>	Reacts with all species
<b>Tested samples</b>	Whole mounts, cell culture, tissue sections
<b>Staining procedure</b>	<a href="#">STAINperfect immunostaining kit A</a>
<b>Format</b>	25µL (approx. 20 tissue sections)

## INFORMATIONS

### Product overview

---

<b>Product name</b>	Serotonin antibody (5-HT rabbit polyclonal Ab) Anti-5-HT antibody
<b>Synonyms</b>	Anti-5-hydroxytryptamine antibody
<b>Immunogen</b>	Conjugated 5-HT When tested in competitive ELISA, the anti-conjugated Serotonin antibody did not show any significant cross reactivity with 5-Hydroxytryptamine analogs, including Tryptamine and 5-Methoxytryptamine
<b>Specificity</b>	
<b>Volume</b>	25µL

### Storage

---

<b>Form</b>	Liquid
<b>Purity</b>	Purified anti-serum
<b>Storage</b>	Store at +4°C for short term (1-2 months). Aliquot and store at -20°C for long term. Avoid repeated freeze / thaw cycles

## PROTOCOLS

**IF – Cell cultures, Whole mounts,  
Tissue sections**

Dilute antibody with the antibody diluent provided in the [STAINperfect immunostaining kit A](#). Use at 1/250 -1/1000 dilution. Follow the STAINperfect protocol suited to your sample

**Comments**

Optimal working dilutions must be determined by the end-user

**Restrictions**

For research use only

**Full protocol**

[Download STAINperfect protocol for Serotonin staining](#)

**Protocols-at-a-glance**

Image not found or type unknown




Image not found or type unknown



[Complete  
Instructions for Use](#)

[Protocol-at-a-glance  
for cell cultures](#)

# REFERENCES

Antibody not yet cited. Submit an article and get a 10% discount.

## Selected papers about serotonin:

- [Gershon MD. 5-Hydroxytryptamine \(serotonin\) in the gastrointestinal tract. Curr Opin Endocrinol Diabetes Obes. 2013 Feb;20\(1\):14-21. doi: 10.1097/MED.0b013e32835bc703.](#)
- [Baganz NL, Blakely RD. A dialogue between the immune system and brain, spoken in the language of serotonin. ACS Chem Neurosci. 2013 Jan 16;4\(1\):48-63. doi: 10.1021/cn300186b. Epub 2012 Dec 7.](#)
- [Bortolato M et al. The role of the serotonergic system at the interface of aggression and suicide. Neuroscience. 2013 Apr 16;236:160-85.](#)
- [Kiyasova V et al. A genetically defined morphologically and functionally unique subset of 5-HT neurons in the mouse raphe nuclei. J Neurosci. 2011 Feb 23;31\(8\):2756-68. doi: 10.1523/JNEUROSCI.4080-10.2011.](#)

## Product pictures

#TITRE

Presence of serotonin in neurons of crayfish brain  
High magnification of serotonergic neurons in the brain of a crayfish. This staining shows serotonin presence in soma containing vesicles and projections. Tissue was processed following whole mount protocol of STAINperfect immunostaining kit A. Fluorescent labeled secondary antibody was used and picture was acquired by confocal imaging.

Image not found or type unknown

#TITRE

Affinity & specificity of anti-5-HT antibody  
Competitive ELISA demonstrates that low amounts of Serotonin conjugate are required to abolish antigen-antibody reaction (high affinity), while rising concentrations of Tryptamine and 5-Methoxy-Tryptamine conjugates do not affect reaction (high specificity).

Image not found or type unknown

#TITRE

Immunostaining of 5-HT-containing cells in adult mouse brain slice  
Anti-Serotonin antibody highlights the soma of serotonergic neurons in the pons of mouse adult brain sagittal section. Staining was performed with STAINperfect immunostaining kit A, following the protocol for tissue sections. Secondary antibody (goat anti-rabbit Alexa Fluor® 488) was used and picture was acquired by confocal imaging at high magnification.

Image not found or type unknown

#TITRE

Staining of serotonergic neurons in mouse primary hindbrain culture  
Anti-Serotonin (green) and MAP2 (red) antibodies permit the detection of serotonergic neurons in primary mouse hindbrain cell culture. Staining was performed using STAINperfect immunostaining kit A, according to the protocol optimized for cell culture and compatible with a MAP2 staining. Fluorescent labeled secondary antibody were used and picture was acquired by high-content imaging.

Image not found or type unknown

#TITRE

Serotonin imaging in embryonic mouse medulla  
Using STAINperfect immunostaining kit A, the anti-Serotonin rabbit polyclonal antibody (Ref: IS1007) demonstrates the presence of serotonin in embryonic mouse medulla (E13.5). The staining was performed according to the protocol optimized for whole-mount samples and was revealed using Alexa Fluor® 488-conjugated secondary antibody.

Image not found or type unknown

#TITRE

Serotonin - 5-hydroxytryptamine (5-HT)  
Serotonin is a monoamine neurotransmitter derived from tryptophan metabolism. Either produced by gut cells or synthesized by serotonergic neurons in the brain, Serotonin is known to be involved in the regulation of appetite, mood, sleep, memory, aging, and to modulate both innate and adaptive immune responses.

Image not found or type unknown

## Contact information

Immusmol  
229 Cours de l'Argonne  
33 000 Bordeaux - France  
Tel: +33 (0) 5 6431 1170  
[www.immusmol.com](http://www.immusmol.com)

**To order, review, ask for technical support, visit product page at:**

<https://www.immusmol.com/shop/serotonin-polyclonal-antibody-bundle/>