

# Private: Epinephrine – Norepinephrine ELISA kit – Research Plasma & Urine samples – FORMER

Ref: BA-E-6500R-F

This Epinephrine/Norepinephrine ELISA kit is tailored for the rapid measurement of Adrenaline (Epinephrine) and Noradrenaline (Norepinephrine) in urine and plasma samples. Intended for research use only, this fast-track quantitative immunoassay enables to determine Epinephrine & Norepinephrine levels in up to 40 samples within 4 hours.

<b>Sample type</b>	10µl urine, 300µl plasma
<b>Samples/kit</b>	40 samples in duplicate
<b>Sensitivity</b>	Epinephrine: 0.3ng/ml urine, 10pg/ml plasma Norepinephrine: 1.5ng/ml urine, 50pg/ml plasma
<b>Range</b>	Epinephrine: 1 - 200ng/ml Norepinephrine: 5 - 1 000ng/ml
<b>Assay time</b>	Sample extraction and acylation (65 min), ELISA (3h)
<b>Reactivity</b>	Reacts with all species
<b>Citations</b>	Not yet cited

# INFORMATIONS

## Product overview

<b>Product name</b>	Epinephrine-Norepinephrine ELISA kit – Urine/Plasma – Fast-Track
<b>Description</b>	Competitive ELISA kit for the quantitative measurement of Adrenaline and Noradrenaline levels in plasma and urine samples. <b>For research use only</b>
<b>Labels</b>	RUO
<b>Format</b>	2 x 96-well plate
<b>Samples</b>	Urine, Plasma
<b>Sample volume</b>	10µl urine, 300µl plasma
<b>Reactivity</b>	Reacts with all species
<b>Standard range</b>	Epinephrine: 0 / 1 – 200ng/ml Norepinephrine: 0 / 5 – 1 000ng/ml
<b>Sensitivity</b>	Epinephrine: 0.3ng/ml urine, 10pg/ml plasma; Norepinephrine: 1.5ng/ml urine, 50pg/ml plasma
<b>Specificity</b>	No significant cross-reactivity was observed with Epinephrine and Norepinephrine analogs such as Dopamine, Metanephrine, Normetanephrine, 3-Methoxytyramine, 3-Methoxy-4-hydroxyphenylglycol, Tyramine, Phenylalanine, Caffeinic acid, L-Dopa, Homovanillic acid, Tyrosine and 3-Methoxy-4-hydroxymandelic acid
<b>Assay time</b>	4 hours
<b>Storage</b>	Store at 2-8°C for to 6 months

**Datasheets**

[Instructions for use](#), [Material safety datasheet](#)

The kit for in-vitro-diagnostic (IVD) use can be ordered with the catalog no. [BA E-6500](#).

## PROTOCOLS

---

**Sample collection & storage**

Plasma: EDTA-Plasma samples only. Do not use haemolytic or lipemic samples. Storage: up to 6 hours at 2 – 8°C; for longer periods (up to 6 months) at – 20°C

Urine: Spontaneous or 24-hours urine, collected in a bottle containing 10-15 mL of 6 M HCl. Storage: for longer periods (up to 6 months) at -20°C

---

**Sample preparation**

Sample preparation, extraction and acylation (65min)

---

**ELISA**

Epinephrine/Norepinephrine antiserum incubation (2H), revelation and read steps (1h)

---

**Detailed protocol**

[Download instructions for use](#)


## REFERENCES

---

Not cited yet. Submit an article and [get a 10% discount](#).


## Product pictures

---



**Fast Epinephrine-Norepinephrine ELISA kit for plasma and urine samples**

---



### **Fast Adrenaline-Noradrenaline ELISA: Standard curves**

Standard curves - to be generated every time the assay is performed.

---

## Contact information

Immusmol  
229 Cours de l'Argonne  
33 000 Bordeaux - France  
Tel: +33 (0) 5 6431 1170  
[www.immusmol.com](http://www.immusmol.com)

**To order, review, ask for technical support, visit product page at:**

[https://www.immusmol.com/?post\\_type=product&p=887](https://www.immusmol.com/?post_type=product&p=887)