

# Private: Epinephrine ELISA kit – Research Plasma & Urine samples – Former

Ref: BA-E-6100R-F

This Epinephrine ELISA kit is tailored for the rapid measurement of Adrenaline (Epinephrine) in urine and plasma samples. Intended for research use only, this fast-track quantitative immunoassay enables to determine Epinephrine levels in up to 40 samples within 4 hours.

<b>Sample type</b>	10µl urine, 300µl plasma
<b>Samples/kit</b>	40 samples in duplicate
<b>Sensitivity</b>	0.3ng/ml urine, 10pg/ml plasma
<b>Range</b>	1 - 200ng/ml
<b>Assay time</b>	Sample extraction and acylation (65 min), ELISA (3h)
<b>Reactivity</b>	Reacts with all species
<b>Citations</b>	Not yet cited

# INFORMATIONS

## Product overview

<b>Product name</b>	Epinephrine ELISA kit – Urine/Plasma – Fast-Track
<b>Description</b>	Competitive ELISA kit for the quantitative measurement of Adrenaline levels in urine and plasma samples. <b>For research use only</b>
<b>Labels</b>	RUO
<b>Format</b>	96-well plate
<b>Samples</b>	10µl urine, 300µl plasma
<b>Reactivity</b>	Reacts with all species
<b>Standard range</b>	0 / 1 – 200ng/ml
<b>Sensitivity</b>	0.3ng/ml urine, 10pg/ml plasma
<b>Specificity</b>	No significant cross-reactivity was observed with Epinephrine analogs such as Dopamine, Noradrenaline, Metanephrine, Normetanephrine, Methoxytyramine, 3-Methoxy-4-hydroxyphenylglycol, Tyramine, Phenylalanine, Caffeinic acid, L-Dopa, Homovanillic acid, Tyrosine and 3-Methoxy-4-hydroxymandelic acid
<b>Assay time</b>	4 hours
<b>Storage</b>	Store at 2-8°C for up to 6 months
<b>Datasheets</b>	<a href="#">Instructions for use</a> , <a href="#">Material safety datasheet</a>

The kit for in-vitro-diagnostic (IVD) use can be ordered with the catalog no. [BA E-6100](#).

## PROTOCOLS

### Sample collection & storage

Plasma: EDTA-Plasma samples only. Do not use haemolytic or lipemic samples. Storage: up to 6 hours at 2 – 8°C; for longer periods (up to 6 months) at – 20°C

Urine: Spontaneous or 24-hours urine, collected in a bottle containing 10-15 mL of 6 M HCl. Storage: for longer periods (up to 6 months) at -20°C

### Sample preparation

Sample preparation, extraction and acylation (65min)

### ELISA

Epinephrine antiserum incubation (2H), revelation and read steps (1h)

### Detailed protocol

[Download instructions for use](#)


## REFERENCES

Not cited yet. Submit an article and [get a 10% discount](#).

## Product pictures



**Fast Urine/Plasma Epinephrine ELISA**



**Fast Adrenaline ELISA: Standard curve**

Standard curve - to be generated every time the assay is performed.

## Contact information

---

Immusmol

229 Cours de l'Argonne

33 000 Bordeaux - France

Tel: +33 (0) 5 6431 1170

[www.immusmol.com](http://www.immusmol.com)

**To order, review, ask for technical support, visit product page at:**

[https://www.immusmol.com/?post\\_type=product&p=892](https://www.immusmol.com/?post_type=product&p=892)