

# Private: Norepinephrine ELISA kit – Research Plasma & Urine samples FORMER

Ref: BA-E-6200R-F

This Norepinephrine ELISA kit is tailored for the rapid measurement of Noradrenaline (Norepinephrine) in urine and plasma samples. Intended for research use only, this fast-track quantitative immunoassay enables to determine Norepinephrine levels in up to 40 samples within 4 hours.

Sample type	10μl urine, 300μl plasma
Samples/kit	40 samples in duplicate
Sensitivity	1.5ng/ml urine, 50pg/ml plasma
Range	5 - 1000ng/ml
Assay time	Sample extraction and acylation (65 min), ELISA (3h)
Reactivity	Reacts with all species
Citations	Not yet cited



### **INFORMATIONS**

<b>Product overview</b>	
Product name	Norepinephrine ELISA kit – Urine/Plasma – Fast-Track
Description	Competitive ELISA kit for the quantitative measurement ofNoradrenaline levels in plasma and urine samples. <b>For research use only</b>
Labels	RUO
Format	96-well plate
Samples	10μl urine, 300μl plasma
Reactivity	Reacts with all species
Standard range	0 / 5 - 1000ng/ml
Sensitivity	1.5ng/ml urine, 50pg/ml plasma
Specificity	No significant cross-reactivity was observed with Noradrenaline analogs such as Dopamine, Adrenaline, Metanephrine, Normetanephrine, Methoxytyramine, 3-Methoxy-4-hydroxyphenylglycol, Tyramine, Phenylalanine, Caffeinic acid, L-Dopa, Homovanillic acid, Tyrosine and 3-Methoxy-4-hydroxymandelic acid
Assay time	4 hours
Storage	Store at 2-8°C for up to 6 months
Datasheets	Instructions for use, Material safety datasheet



The kit for in-vitro-diagnostic (IVD) use can be ordered with the catalog no. BA E-6200.

#### **PROTOCOLS**

Sample collection & storage	<u>Plasma:</u> EDTA-Plasma samples only. Do not use haemolytic or lipemic samples. Storage: up to 6 hours at 2 – 8°C; for longer periods (up to 6 months) at – 20°C <u>Urine:</u> Spontaneous or 24-hours urine, collected in a bottle containing 10-15 mL of 6 M HCl. Storage: for longer periods (up to 6 months) at -20°C
Sample preparation	Sample preparation, extraction and acylation (65min)
ELISA	Norepinephrine antiserum incubation (2H), revelation and read steps (1h)
Detailed protocol	Download instructions for use

#### **REFERENCES**


Not cited yet. Submit an article and get a 10% discount.

## **Product pictures**



Plasma Norepinephrine ELISA kit





#### Plasma Noradrenaline ELISA: Standard curve

Standard curve - to be generated every time the assay is performed.

#### Contact information

Immusmol 229 Cours de l'Argonne 33 000 Bordeaux - France Tel: +33 (0) 5 6431 1170

www.immusmol.com

To order, review, ask for technical support, visit product page at:

https://www.immusmol.com/?post\_type=product&p=911