

BA-E-6200

Fast Norepinephrine ELISA kit - Plasma & Urine

This fast Norepinephrine ELISA kit is tailored to rapidly measure Noradrenaline (Norepinephrine) in urine and plasma samples. Intended for *in vitro* diagnostics & research use, this quantitative immunoassay enables the determination of Norepinephrine levels in up to 40 samples within 4 hours.

[Cited in 8 papers](#)



Sample type	10µL urine, 300µL plasma
Capacity	96-well plate
Sensitivity	Urine: 1.5 ng/mL Plasma: 50 pg/mL
Range	5 - 1 000 ng/mL
Assay time	4h
Reactivity	Reacts with all species
Application	<i>In vitro</i> diagnostics (EU only), or Research use
Storage	Store at 2-8°C for to 6 months
Sample collection & storage	Plasma: EDTA-Plasma samples only. Do not use haemolytic or lipemic samples. Storage: up to 6 hours at 2 - 8°C; for longer periods (up to 6 months) at - 20°C Urine: Spontaneous or 24-hour urine, collected in a bottle containing 10-15 mL of 6 M HCl. Storage: for longer periods (up to 6 months) at -20°C
Sample preparation	Sample preparation, extraction and acylation (65min)
ELISA	Norepinephrine antiserum incubation (2h), revelation and read steps (1h)

To see data, publications, and download instructions for use and MSDS, visit:
<https://www.immusmol.com/product/fast-norepinephrine-elisa-kit/>