IS010

Kynurenic acid Antibody - Mouse Monoclonal

The monoclonal IS010 anti-Kynurenic acid antibody was validated for IHC and IF in human caudate putamen tissues. Competitive ELISA demonstrated the antibody to be highly affine and specific.
Cited in 1 paper

Clonality	Monoclonal antibody (clone 4G12-A12)
Host	Mouse
Reactivity	Reacts with all species
Tested samples	Human brain tissue sections
Staining procedure	Perform heat antigen retrieval (pH=6) before standard IHF/IF staining
Format	50µL (approx. 40 tissue sections)

Storage	Store at +4°C for short term (6 months). Aliquot and store at -20°C for long term. Avoid repeated freeze / thaw cycles
Immunohistochemistry (IHC)	Dilute at 1:200-1:2000. Perform heat antigen retrieval (pH=6) before initiating IHC staining protocol on paraffin-embedded and frozen sections
Immunofluorescence (IF)	Dilute at 1:100-1:1000 on paraffin-embedded and frozen sections. Perform heat antigen retrieval and incubate with fluorescent dyes conjugated secondary antibody
Comments	Optimal working dilutions must be determined by the end-user
Restrictions	For research use only

To see data, publications, and download instructions for use and MSDS, visit:
https://www.immusmol.com/product/kynurenic-acid-mab/

Immusmol, 229 Cours de l'Argonne, 33000 Bordeaux – France.
Email : contact@immusmol.com | Tel : +33 (0)5 64 31 11 70 | www.immusmol.com